

- 77 **Bailey DB Jr**, Mesibov GB, Hatton DD, Clark RD, Roberts JE, Mayhew L. Autistic behavior in young boys with fragile X syndrome. *J Autism Dev Disord* 1998;**28**:499-508.
- 78 **Fisch GS**. Is autism associated with the fragile X syndrome? *Am J Med Genet* 1992;**43**:47-55.
- 79 **Hallmayer J**, Pintado E, Lotspeich L, Spiker D, McMahon W, Petersen PB, Nicholas P, Pingree C, Kraemer HC, Wong, DL. Molecular analysis and test of linkage between the FMR-1 gene and infantile autism in multiplex families. *Am J Hum Genet* 1994;**55**:951-9.
- 80 **Lam CW**, Yeung WL, Ko CH, Poon PM, Tong SF, Chan KY, Lo IF, Chan LY, Hui J, Wong V, Pang CP, Lo YM, Fok TF. Spectrum of mutations in the MECP2 gene in patients with infantile autism and Rett syndrome. *J Med Genet* 2000;**37**:e41.
- 81 **Khan SG**, Levy HL, Legerski R, Quackenbush E, Reardon JT, Emmert S, Sancar A, Li L, Schneider TD, Cleaver JE, Kraemer KH. Xeroderma pigmentosum group C splice mutation associated with autism and hypoglycinemia. *J Invest Dermatol* 1998;**111**:791-6.
- 82 **Gorlin RJ**, Cohen MM, Hennekam RC. *Syndromes of the head and neck*. 4th ed. New York: Oxford University Press, 2001.
- 83 **Lowe T**, Tanaka K, Seashore MR, Young JG, Cohen DJ. Detection of phenylketonuria in autistic and psychotic children. *JAMA* 1980;**243**:126-8.
- 84 **Jaeken J**, Van den Berghe G. An infantile autistic syndrome characterised by the presence of succinylpurines in body fluids. *Lancet* 1984;**ii**:1058-61.
- 85 **Naruse H**, Hayashi T, Takesada M, Nakane A, Yamazaki K. Metabolic changes in aromatic amino acids and monoamines in infantile autism and development of new treatment related to the finding. *No To Hattatsu* 1989;**21**:181-9.
- 86 **Coleman M**, Blass JP. Autism and lactic acidosis. *J Autism Dev Disord* 1985;**15**:1-8.
- 87 **Kelley R**. Inborn errors of cholesterol biosynthesis. *Adv Pediatr* 2000;**47**:1-53.
- 88 **Dennis M**, Lockyer L, Lazenby AL, Donnelly RE, Wilkinson M, Schoonheydt W. Intelligence patterns among children with high-functioning autism, phenylketonuria, and childhood head injury. *J Autism Dev Disord* 1999;**29**:15-17.
- 89 **Seim A**, Reichelt KL. An enzyme/brain-barrier theory of psychiatric pathogenesis: unifying observations on phenylketonuria, autism, schizophrenia and post-partum psychosis. *Med Hypotheses* 1995;**45**:498-502.
- 90 **Shaw W**, Kassen E, Chaves E. Increased urinary excretion of analogs of Krebs cycle metabolites and arabinose in two brothers with autistic features. *Clin Chem* 1995;**44**:1094-104.
- 91 **Lombard J**. Autism: a mitochondrial disorder? *Med Hypotheses* 1998;**50**:497-500.
- 92 **Jaeken J**, Matthijs G, Barone R, Carchon H. Carbohydrate deficient glycoprotein (CDG) syndrome type I. *J Med Genet* 1997;**34**:73-6.
- 93 **de Lonlay P**, Seta N, Barrot S, Chabrol B, Drouin V, Gabriel BM, Journel H, Kretz M, Laurent J, Le Merrer M, Leroy A, Pedespan D, Sarda P, Villeneuve N, Schmitz J, van Schaftingen E, Matthijs G, Jaeken J, Korner C, Munnich A, Saudubray J M, Cormier-Daire V. A broad spectrum of clinical presentations in congenital disorders of glycosylation I: a series of 26 cases. *J Med Genet* 2001;**38**:14-19.
- 94 **White P**, DeMeyer W, DeMeyer M. EEG abnormalities in early childhood schizophrenia. *Am J Psychiatry* 1964;**120**:1950-8.
- 95 **Minshew NJ**. Indices of neural function in autism: clinical and biologic implications. *Pediatrics* 1991;**87**:774-80.
- 96 **Courchesne E**. Neuroanatomic imaging in autism. *Pediatrics* 1991;**87**:781-90.
- 97 **Courchesne E**. An MRI study of autism: the cerebellum revisited. *Neurology* 1999;**52**:1106-7.
- 98 **Hennekam RC**. Abnormal mental development. In: Tindall B, ed. *Balliere's clinical paediatrics*. London: Balliere Tindall 1998:317-32.
- 99 **Risch N**, Spiker D, Lotspeich L, Nouri N, Hinds D, Hallmayer J, Kalaydjieva L, McCague P, Dimiceli S, Pitts T, Nguyen L, Yang J, Harper C, Thorpe D, Vermeer S, Young H, Hebert J, Lin A, Ferguson J, Chiotti C, Wiese-Slater S, Rogers T, Salmon B, Nicholas P, Myers RM. A genomic screen of autism: evidence for a multilocus etiology. *Am J Hum Genet* 1999;**65**:493-507.

Familial pericentric inversion of chromosome 5 in a family with benign neonatal convulsions

D Concolino, M A Iembo, E Rossi, S Giglio, G Coppola, E Miraglia del Giudice, P Strisciuglio

J Med Genet 2002;**39**:214-216

We describe a family in whom a pericentric inversion of chromosome 5 segregates with benign familial neonatal convulsions (BFNC). BFNC is an autosomal dominant form of epilepsy characterised by spontaneous partial or generalised clonic convulsions beginning within the first months of life. Seizures usually disappear by the age of 6 months; intercritical electroencephalogram and subsequent psychomotor development are normal. BFNC loci have been mapped to human chromosomes 20q13.3 (*BFNC1*) and 8q24 (*BFNC2*), based on linkage analysis.^{1,2} Recently, two potassium channel genes (*KCNQ2* and *KCNQ3*), located in these two regions, were shown to be mutated in *BFNC1* and *BFNC2*, respectively.^{3,4}

We report a family with BFNC and a pericentric inversion of chromosome 5 cosegregating with BFNC. Fluorescence in situ hybridisation (FISH) experiments were performed to define the breakpoints at YAC level. The linkage of BFNC to *KCNQ2* and mutations in the *KCNQ3* gene were excluded. This report raises the possibility of a new locus for BFNC on chromosome 5.

CASE REPORT

The proband, a male, was the second child of unrelated parents, born at 40 weeks of gestation after an uneventful pregnancy. Apgar score was 10 at one and five minutes. The

parents were both 32 years old at his birth. Birth weight was 3400 g (50-75th centile), length was 51 cm (50-75th centile), head circumference was 35 cm (50-75th centile), and the clinical examination was normal except for the presence of hypotelorism. On the third day of life he had five episodes of generalised clonic convulsions treated with phenobarbital (5 mg/kg/day). Two additional episodes of seizures occurred during the first months of life and therefore long term therapy was continued. Routine laboratory investigations including plasma ammonia levels and acid base status were normal. Plasma and urinary amino acids were normal. Brain CT scan, electroencephalogram, echocardiography, and abdominal echography were all normal. At 8 months of age he was developmentally normal and continued to take phenobarbital.

The mother and maternal grandmother appeared to be intellectually normal but reported a clear history of convulsions in infancy and afebrile seizures later in childhood. Both showed hypotelorism without other clinical or dysmorphic signs. No other relatives of the proband had hypotelorism or a history of idiopathic neonatal convulsions.

Cytogenetic and molecular analysis

Chromosome analysis was performed on QFQ and GTG banded metaphases from synchronised peripheral lymphocyte cultures using standard procedures. The proband's




Figure 1 (A) Ideograms of the normal and inverted chromosome. (B) GTG banded partial karyotype of the patient. The normal chromosome 5 is on left and the inverted one on the right.

karyotype was interpreted as 46,XY,inv(5)(p15q11) (fig 1) and the same inversion was present in the proband's mother and maternal grandmother but not in other relatives (fig 2).

Inversion breakpoints were defined at the YAC by fluorescence in situ hybridisation (FISH) to chromosome preparations from the proband. YACs containing chromosome 5 specific sequences from several locations on the p and q arms were selected according to the Genome Database (www.genome.wi.mit.edu/cgi-bin/conting/phys_map). YAC DNA was labelled with biotin using nick translation. The labelled probes were visualised with FITC-avidin (Vector) and the chromosomes were counterstained with DAPI (Sigma). Hybridisations were analysed with a Zeiss Axioplan epifluorescence microscope and images were captured with the Power Gene FISH System (PSI).

Breakpoints were found at YACs 956a11 (5p15.1, D5S1954-D5S416 at 28 cM from the short arm telomere) and 854b12 (5q11.2, D5S2076-D5S664 at 63 cM) (fig 3). Therefore the full karyotype of the proband was 46,XY,inv(5)(p15q11).ish inv(5)(p15.1q11.2)(D5S1954sp,D52076sp).

Segregation of the polymorphic markers D8S1385 and D8S558 closely linked to *KCNQ3* and of the Thr752Asn polymorphism located in exon 16 of the *KCNQ2* gene was studied in the family members. This study excluded linkage of BFNC to *KCNQ2* while linkage to *KCNQ3* was not excluded. Therefore, the *KCNQ3* coding region, including the exon-intron boundaries, was analysed in the patient and his mother. Polymerase chain reactions (PCR) were performed using primers and PCR conditions kindly provided by Shinichi Hisose, Department of Paediatrics, Fukouka University, Japan. PCR products were directly sequenced using ABI PRISM 310 sequencer and the Big Dye Terminator Cycle Sequencing Ready Reaction Kit (Applied Biosystem, Foster City, CA). No mutations were found.




Figure 2 Family pedigree.

DISCUSSION

This report is another example of a rare case of an inherited pericentric inversion of chromosome 5 with the presence of clinical findings. Other pericentric inversions of the same region of chromosome 5 have been described,^{5,6} but not with the same breakpoints in both arms. In our family, FISH analysis localised the 5p breakpoint at YAC clone 956a11 located in proximal 5p15.1 and the 5q breakpoint at YAC 854b12 located in 5q11.2. The breakpoint in the short arm of chromosome 5 in this report is not in the cri du chat syndrome critical region, mapped in 5p15.3 for high pitched cry and in 5p15.2 for the remaining features.⁷ Therefore, this is the first example of a family with a pericentric inversion of chromosome 5 and BFNC.

In BFNC, seizures begin in the neonatal period and generally have a favourable outcome with spontaneous remission in the first year of life. The clinical heterogeneity of this disorder has been suggested by differences among pedigrees in the risk of developing epilepsy in later life.⁸ In the last few years, two loci for BFNC have been mapped to chromosomes 20q13.3 and 8q24^{1,2} showing that this disorder is also genetically heterogeneous. The majority of families are linked to chromosome 20q with mutations in the potassium channel gene *KCNQ2*. Only two families have been detected with mutations in the highly homologous gene *KCNQ3* at 8q24,^{9,10} so *KCNQ2* was considered to be a major gene locus for BFNC.^{3,4,9} Mutations in both genes are heterozygous, according to the autosomal dominant inheritance pattern of BFNC. A new benign idiopathic epilepsy in early life, with a favourable outcome and autosomal dominant inheritance, has been reported.¹¹ In this family, the seizures began between 4 and 7 months of life and for this reason the condition was called benign infantile familial convulsions (BFIC). A locus for BFIC has been mapped to chromosome 19q12-13,¹² suggesting that BFNC and BFIC are not allelic diseases with variable expressivity but are different entities. However, in the seven families with BFIC recently investigated, no evidence of linkage with chromosome 19 markers has been found.¹³

In conclusion, there are several lines of evidence for clinical and genetic heterogeneity of this condition and at least three loci responsible for autosomal dominant benign epilepsy of early life. Our family shows a strong correlation between the chromosome 5 inversion and BFNC. In fact, the disease was present only in the three inv(5) subjects but not in the two




Figure 3 FISH analysis with the YAC 956a11 (left) and YAC 854b12 (right) crossing the inv(5) breakpoints on the short and long arm, respectively.

aunts of the proband or in his brother. We hypothesised that breakage or transcription silencing of a gene at one of the inversion breakpoints is responsible for the disease. Moreover, the exclusion of linkage of BFNC to *KCNQ2* and *KCNQ3* in this family supports this hypothesis.

Authors' affiliations

D Concolino, M A Iembo, P Strisciuglio, Department of Paediatrics, University of Catanzaro "Magna Graecia", Italy
E Rossi, S Giglio, Department of General Biology and Medical Genetics, University of Pavia, Italy
G Coppola, E Miraglia del Giudice, Department of Paediatrics, 2nd University of Naples, Italy

Correspondence to: Dr P Strisciuglio, Department of Paediatrics, Faculty of Medicine, University of Catanzaro, C/o Ospedale "A Pugliese", Viale Pio X, 88100 Catanzaro, Italy; Strisciuglio-unicz@libero.it

REFERENCES

- 1 Leppert M, Anderson VE, Quattlebaum T, Stauffe D, O'Connell P, Nakamura Y, Lalouel JM, White R. Benign familial neonatal convulsions linked to genetic markers on chromosome 20. *Nature* 1989;**337**:647-8.
- 2 Lewis TB, Leach RJ, Ward K, O'Connell P, Ryan SG. Genetic heterogeneity in benign familial neonatal convulsions: identification of a new locus on chromosome 8. *Am J Hum Genet* 1993;**53**:670-5.
- 3 Biervet C, Schroeder BC, Kubisch C, Berkovic SF, Propping P, Jentsch TJ, Steinlein OK. A potassium channel mutation in neonatal human epilepsy. *Science* 1998;**279**:403-6.
- 4 Singh NA, Charles C, Stauffer D, DuPont BR, Leach RJ, Melis R, Ronen GB, Bjerre I, Quattlebaum T, Murphy JV, McHarg ML, Gagnon D, Rosales TO, Peiffer A, Anderson VE, Leppert M. A novel potassium channel gene, *KCNQ2*, is mutated in inherited epilepsy of newborns. *Nat Genet* 1998;**18**:25-9.
- 5 Beemer FA, de France HF, Rosina Angelista IJ, Cast BP, Guyt R. Familial partial monosomy 5p and trisomy 5q; three cases due to paternal pericentric inversion 5 (p151q333). *Clin Genet* 1984;**26**:209-15.
- 6 Goodart SA, Butler MG, Overhauser J. Familial double pericentric inversion of chromosome 5 with some features of cri-du-chat syndrome. *Hum Genet* 1996;**97**:802-7.
- 7 Overhauser J, Huang X, Gersh M, Wilson W, McMahon J, Bengtsson U, Rojas K, Meyer M, Wasmuth JJ. Molecular and phenotypic mapping of the short arm of chromosome 5: sublocalization of the critical region of the cri-du-chat syndrome. *Hum Mol Genet* 1994;**3**:247-52.
- 8 Shevell MI, Sinclair DB, Metrakos K. Benign familial neonatal seizures: clinical and electroencephalographic characteristics. *Pediatr Neurol* 1986;**2**:272-5.
- 9 Charlier C, Singh NA, Ryan SG, Lewis TB, Reus BE, Leach RJ, Leppert M. A pore mutation in a novel KQT-like potassium channel gene in a idiopathic epilepsy family. *Nat Genet* 1998;**18**:53-5.
- 10 Hirose S, Zenri F, Akiyoshi H, Fukuma G, Iwata H, Inoue T, Yonetani M, Tsutsumi M, Muranaka H, Kurokawa T, Hanai T, Wada K, Kaneko S, Mitsudome A. A novel mutation of *KCNQ3* (c.925T→C) in a Japanese family with benign familial neonatal convulsions. *Ann Neurol* 2000;**47**:822-6.
- 11 Vigeveno F, Fusco L, Di Capua M, Ricci S, Sebastianelli R, Lucchini P. Benign infantile familial convulsions. *Eur J Pediatr* 1992;**151**:608-12.
- 12 Guipponi M, Rivier F, Vigeveno F, Beck C, Crespel A, Echenne B, Lucchini P, Sebastianelli R, Baldy-Moulinier M, Malafosse A. Linkage mapping of benign familial infantile convulsions to chromosome 19q. *Hum Mol Genet* 1997;**6**:473-7.
- 13 Gennaro E, Malacarne M, Carbone I, Riggio MC, Bianchi A, Bonanni P, Boniver C, Dalla Bernardina B, De Marco P, Giordano L, Guerrini R, Santorum E, Sebastianelli R, Vecchi M, Vegiotti P, Vigeveno F, Bricarelli FD, Zara F. No evidence of a locus for benign familial infantile convulsions on chromosome 19q12-13. *Epilepsia* 1999;**40**:1799-803.